

**EVALUATION OF PUMPKIN CULTIVARS  
FOR SOUTHERN OHIO, 2008**

Brad R. Bergefurd, Dr. Shawn Wright, Thom Harker,  
Wayne Lewis, Al Welch, Lynn Miller  
The Ohio State University South Centers  
1864 Shyville Road, Piketon, Ohio 45661-9749  
Phone: (740) 289-2071

**The authors wish to thank the Ohio Vegetable and Small Fruit Research and Development Program and cooperating seed companies for providing funding and support of this project.**

**METHODS:**

Seeds were planted June 19<sup>th</sup>. Rows were 10 foot apart with seeds planted 3' apart in the row. Experimental design was randomized complete block with 4 replications. The field is located in southern Ohio, at the Ohio State University South Centers research and demonstration plots. 120 units of N were applied prior to planting. A standard commercial fungicide and insecticide program following OSU Bulletin #672, The Ohio Vegetable Production Guide, was followed, on a 7-10 day schedule.

**RESULTS:**

Three varieties were tested in replicated trials including Gladiator, Spartan and Camaro. Camaro had the lowest number of orange fruit per acre (1000) and the lowest pounds of orange fruit per acre (16,104). Average fruit size for all entries ranged between 15 and 19 pounds.

Ten varieties were planted as observation trials only by seed company request including Treasure, Progress, Packer, Capital, Superior, Summit, Warlock, Magic Wand, Field Trip and Gargoyle.

**Table 1: Orange fruit per acre.**

<b>Cultivar</b>	<b>Marketable fruit</b>
Gladiator	2,214
Spartan	1,928
Camaro	1,000

**LSD 868**

**Table 2: Orange pounds per acre.**

<b>Culivar</b>	<b>Marketable pounds</b>
Gladiator	34,271
Spartan	33,382
Camaro	16,104

**LSD 14041**